



## Safer Garden Fungicide - Concentrate

### For Flowers, Fruit & Vegetables

*ALWAYS SHAKE WELL BEFORE USING*

#### **PRECAUTIONARY HAZARDS:**

##### **Hazards to Humans and Domestic Animals:**

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

**FIRST AID: If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment advice.**

##### **ENVIRONMENTAL HAZARDS:**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of rinsate or equipment washwater.

#### **DIRECTIONS FOR USE:**

*It is a violation of Federal law to use this product in a manner inconsistent with its labeling.*

##### ***SHAKE WELL.***

For control of **Black Spot, Leaf Spot** and **Rust**, mix 1 part concentrate to 30 parts water or 4 fl. oz. (8 tbsp.) per gal. of water.

For **Powdery Mildew**, mix 2 fl. oz. (4 tbsp.) per gallon of water. Place about half of the volume of water in sprayer, add concentrate, stir well, add rest of water and mix thoroughly. Run agitator on large tank sprayers.

For coverage, see individual applications. Spray plants thoroughly to wet the upper and lower surfaces of foliage. Repeat as recommended.

To control powdery mildew, leaf spot, and rust on **Flowers, Foliage Plants & Ornamentals**, spray at 7 to 10 day intervals or as needed.

To control powdery mildew, black spot and rust on **Roses**, spray at 7 to 10 day intervals, or following rain. Be sure to treat new growth & buds. Avoid spraying when plants are in full sun or temperature exceeds 85 degrees F. Apply 1 gal. mix to each 350 sq. ft.

To control powdery mildew on **Beans & Peas**, spray at first sign of disease & repeat at 7 day intervals as necessary. May be used up to day before harvest. Apply 1 gal. of mix to each 250 sq. ft.

To control powdery mildew on **Apples & Pears**, apply to thoroughly wet foliage. Apply at pre-bloom stage or after petal fall as needed. **NOTE:** Some apple & pear varieties are sulfur sensitive; test for varietal sensitivity prior to full scale treatment.

To control powdery mildew on **Potatoes**, spray before disease appears & repeat as necessary. Apply 1 gal. of mix to each 350 sq. ft.

On **Cucumbers and Squash**, spray when powdery mildew appears & repeat as necessary. Do not spray young plants. Apply 1 gal. of mix to each 600 sq. ft.

To control powdery mildew on **Grapes**, thoroughly cover foliage. Begin when new shoots are 6-10 inches. Repeat before blooms open and at 14 day intervals as needed. **CAUTION:** Do not apply to wine grapes within 21 days of harvest. Sulfur may damage concord & other labrusca varieties.

To control powdery mildew on **Strawberries**, spray when disease appears and repeat as necessary. Do not spray young plants. Apply 1 gal. of mix to each 300 sq. ft. **NOTE:** Do not apply to strawberries that will be used for canning. May be used up to the day before harvest.

#### **HOME GREENHOUSES:**

To control powdery mildew, leaf spot and rust on bedding plants, propagates, fruits & vegetables: Mix 1 part concentrate to 30 parts water (8 tbsp. per gal.) and apply according to above recommendations by crop type. Test for phytotoxicity before full-scale application.

#### **USE CAUTIONS:**

Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not apply on Boston fern, spinach, apricots, filberts, walnuts, and viburnum. Do not apply when temperature exceeds 85 degrees F. Do not apply for at least 4 weeks

following application of an oil spray or within 4 days of other pesticides. Be sure to wash and rinse spray equipment thoroughly after each use. Do not allow mixed spray to stand in spray equipment. **Re-entry:** Avoid contact with plants until spray has dried.

## **STORAGE AND DISPOSAL:**

**Pesticide Storage:** Store only in original container in a cool, dry area inaccessible to children and pets. Protect from freezing and heat. Keep container tightly closed to prevent evaporation.

**Pesticide Disposal and Container Handling:** Non-refillable container. Do not reuse or refill this container. **If Empty:** Place in trash or offer for recycling if available.

**If Partly Filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Distributed by Safer®, Inc., a wholly-owned subsidiary of Woodstream Corporation.

Woodstream  
69 North Locust Street, Lititz, PA 17543  
1-800-800-1819  
717-626-2125  
EPA Reg. No. 42697-37  
EPA Est. No. 63416-MN-01

Look for other Safer® Brand products at [www.saferbrand.com](http://www.saferbrand.com)

© Copyright 2009 Woodstream Corp. LBL5456B-2 Model No. 5456 Made in USA